

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 174 514 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
23.01.2002 Bulletin 2002/04

(51) Int Cl.7: **C12N 15/86**, A61K 48/00,
C12N 7/04, A61K 39/145

(21) Application number: 00115626.4

(22) Date of filing: 20.07.2000

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(71) Applicant: **ARTEMIS Pharmaceuticals GmbH**
51063 Köln (DE)

(72) Inventors:
• **Hobom, Gerd**
35392 Glessen (DE)

• **Menke, Anette**
35037 Marburg (DE)
• **Meyer-Rogge, Sabine**
35321 Laubach-Münster (DE)

(74) Representative:
Helbing, Jörg, Dr.Dipl.-Chem. et al
Patentanwälte von Kreisler-Selting-Werner,
Deichmannhaus am Hauptbahnhof
50667 Köln (DE)

(54) **Recombinant influenza viruses with bicistronic vRNAs coding for two genes in tandem arrangement**

(57) The invention relates to recombinant influenza viruses for high-yield expression of incorporated foreign gene(s), which are genetically stable in the absence of any helper virus and which comprise at least one viral RNA segment being a tandem bicistronic RNA molecule coding for two genes in tandem, in said tandem bicistronic RNA molecule one of the standard viral genes being in covalent junction with a foreign, recombinant gene and having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding

region.

The invention further provides a method for obtaining attenuated viruses which resist reassortment dependent progeny production in case of chance superinfections by wild-type influenza viruses; a method for the production of said recombinant influenza viruses; pharmaceutical compositions comprising said recombinant influenza viruses; and the use of said recombinant influenza viruses for preparing medicaments for vaccination purposes.

EP 1 174 514 A1

Description**Field of the Invention**

5 [0001] The invention relates to recombinant influenza viruses for high-yield expression of incorporated foreign gene (s), which are genetically stable in the absence of any helper virus and which comprise at least one viral RNA segment being a tandem bicistronic RNA molecule coding for two genes in tandem, in said tandem bicistronic RNA molecule one of the standard viral genes being in covalent junction with a foreign, recombinant gene and having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region. In particular the above
 10 tandem bicistronic RNA molecule contains one of the standard viral genes in distal mRNA position behind a foreign, recombinant gene in proximal position, or vice versa, both in antisense orientation with regard to the viral RNA within the virus. For simultaneous expression of both genes the proximal reading frame is flanked by splice donor and acceptor signals which have the quality to allow a partial yield of spliced mRNA only, i.e., resulting in the presence of both, spliced and unspliced mRNA simultaneously.

15 [0002] The invention further provides a method for obtaining attenuated viruses which resist reassortment dependent progeny production in case of chance superinfections by wild-type influenza viruses; a method for the production of said recombinant influenza viruses; pharmaceutical compositions comprising said recombinant influenza viruses; and the use of said recombinant influenza viruses for preparing medicaments for vaccination purposes.

Technical Background

[0003] Redesigning influenza virus into a vector system for expression of foreign genes similar to what has been achieved in several other thoroughly studied viruses such as adenovirus, retrovirus, Semliki Forest virus or Rabies virus has the advantage of an industrially well established mode of cheap propagation for influenza in fertilized chicken
 25 eggs leading to rather high titers (above 10^{10} /ml). On the other hand none of the constituent vRNA segments may be deleted from the influenza genome according to our present knowledge, and give room for large-size foreign insertions. Only small fragments of foreign polypeptide chains such as B cell epitopes (10 to 15 amino acids) may be inserted into selected positions within two of the viral proteins, i.e. in exchange for one of the variable antigenic regions located at the surface of hemagglutinin (Muster et al., Mucosal model of immunization against human immunodeficiency virus
 30 type 1 with a chimeric influenza virus, J. Virol. 69 (11), 6678-6686 (1995)), or into the stalk sequence of viral neuraminidase (Garcia-Sastre and Palese, The cytoplasmic tail of the neuraminidase protein of influenza A virus does not play an important role in the packaging of this protein into viral envelopes, Virus Res. 37, 37-47 (1995)), and be stably maintained as functional fusion proteins. Constructs of this kind turned out to be useful for experimental vaccination in a few cases studied, but only rather few clearly defined epitope sequences (of ten to twelve amino acids each) are
 35 known today, and some of them might also be misfolded within such restricted fusion protein positions, or in other cases interfere with formation of the correct tertiary structure and function of their host polypeptide chains. Incorporation of a full-size foreign protein into influenza virus via reverse genetics, encoded by an independent ninth vRNA molecule in addition to its regular set of eight standard vRNA segments is without special provisions only transiently possible (Luytjes et al., Amplification, expression, and packaging of a foreign gene by influenza virus. Cell 59,
 40 1107-1113 (1989); Enami et al., An influenza virus containing nine different RNA segments, Virology 185, 291-298 (1991)). In the absence of a continuous selective pressure any additional recombinant vRNA segment cannot be stably maintained as long as the wildtype promoter sequence is used on that ninth vRNA segment, and it will inadvertently be lost after few steps of viral propagation.

Using a different system of influenza reverse genetics developed in our laboratory (Zobel et al., RNA polymerase I
 45 catalysed transcription of insert viral cDNA, Nucleic Acids Res. 21, 3607-3614 (1993); Neumann et al., RNA polymerase I-mediated expression of influenza viral RNA molecules, Virology 202, 477-479 (1994)), which was built around *in vivo* synthesis of recombinant vRNA molecules by cellular RNA polymerase I transcription of the respective template cDNA constructs, modified terminal viral RNA sequences (hereinafter "promoter-up mutations" or promoter-up variants") have been designed by nucleotide substitutions (Neumann and Hobom, Mutational analysis of influenza virus promoter
 50 elements *in vivo*, J. Gen. Virol. 76, 1709-1717 (1995); WO 96/10641). The above promoter-up variants carry up to five nucleotide substitutions (in promoter-up variant 1920; see Flick and Hobom, J. Gen. Virol. 80, 2565-2572 (1999)). When these promoter-up variants are attached to a recombinant ninth vRNA segment its increased transcription and amplification rate will not only compensate for the losses suffered spontaneously, but even cause accumulation of the foreign vRNA segment during simple viral passaging, in the absence of any selection.

55 [0004] However, due to its over-replication relative to all of the regular influenza vRNA segments (which of course are connected to wild-type promoter sequences) after catching up with the others the foreign segment will become over-abundant. This increasingly will result in viral particles that have incorporated several copies of recombinant vRNA, but no longer have a full set of all eight standard segments incorporated among an average of about 12-15 vRNA

molecules present within a virion. Such particles are defective and will not result in plaque formation, hence after an initial increase of recombinant viral particles during the first steps of propagation a dramatic decrease is observed, usually at the third or fourth step of viral passaging, depending on the size of the recombinant vRNA and the level of the promoter-up mutation attached.

[0005] A balanced situation with regard to the insert length and the level of promoter activity can be achieved, and has been propagated in a particular case over 11 passages, with essentially stable levels of recombinant viruses among a majority of helper viruses (around 80%) during these steps. If a full-level promoter-up mutation is used (1104 or the variant 1920, see below) a balanced-level propagation is reached in conjunction with a recombinant vRNA size of 4000 nucleotides (Maysa Azzeh, Ph.D. Thesis, Univ. Giessen (2000)).

In all of these preparations, both in transiently achieved increased yields (up to 40% of recombinants after three or four steps of viral passage), and in a balanced propagation of recombinant influenza viruses (10 - 20%) the respective viral progeny inadvertently constitute mixtures with a majority of non-recombinant helper viruses. These result both from a statistical mode of packaging vRNA molecules into a virion (the ninth segment may not be co-packaged), and from the fraction of cells solely infected by helper virus.

To solve the problems of fractional yields and of instability during viral propagation of recombinant influenza, it was suggested to use a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus and which comprises at least one viral RNA segment being an ambisense RNA molecule (designated "ambisense RNA segment") and containing one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, or *vice versa*, in overall convergent arrangement (PCT/EP00/01903). The ambisense RNA segment preferably should contain the promoter-up mutations. The PCT/EP00/01903 moreover discloses a method of constructing specific influenza carrier (helper) strains carrying one or more ribozyme target sites (of type one) in vRNA flanking positions comprising

(a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, carrying two different 3' promoter sequences in tandem (an external promoter-up variant and an internal wild-type promoter), which are separated by a second type of ribozyme target sequence, and which carry the said internal ribozyme target sites of type one;

(b) followed by infection of an influenza wildtype strain;

(c) thereafter amplification through simple steps of viral propagation; and

(d) finally isolation through removal of their external 3' promoter sequence by ribozyme cleavage through infection of cells expressing ribozyme type 2, followed by plaque purification.

[0006] The resulting special helper virus strains carrying a vRNA segment with external ribozyme target sites of type 1 in exchange for the equivalent regular vRNA molecule are then used for the rescue of ambisense RNA molecules. These are exclusively maintained in the recombinant viruses after passage of viral propagation through ribozyme (type 1) containing host cells, which will destroy the sensitive vRNA molecules of the specially prepared helper viruses.

[0007] However, the above ambisense constructs are susceptible to (intra-nuclear) mRNA double-strand formation, which will partially reduce the expression rates of both the ambisense genes, in particular the gene driven by the (weaker) cRNA promoter. The fluctuating extent of this effect made it difficult to bring the expression rate of the influenza gene within the ambisense segment into balance with other influenza genes.

This was the problem to be solved with the present invention.

Summary of the Invention

[0008] Starting out from two observations in this laboratory which are discussed above and which concern two hitherto unsuspected properties of influenza viral RNA polymerase in its interaction with terminally adapted influenza-specific RNA molecules, stable recombinant influenza viruses were found, which solve the above problems.

[0009] The recombinant viruses of the present invention can be used for cheap propagation in fertilized eggs, either for production of those recombinant viruses themselves or for production of foreign proteins or glycoproteins encoded by them, and hence find application in (glyco)protein production or in providing vector systems for somatic gene therapy or in being used as vaccination agents.

[0010] Thus, the present invention provides

(1) a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus and which comprises at least one viral RNA segment being a bicistronic RNA molecule coding for two genes in tandem arrangement (hereinafter "tandem bicistronic RNA segment" or "tandem RNA segment"), in said tandem RNA segment one of the standard viral genes being in covalent junction with a foreign, recombinant gene and said tandem RNA segment having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region;

(2) a preferred embodiment of the recombinant influenza virus defined in (1) above, in which the terminal viral RNA sequences of said at least one tandem RNA segment, which are active as the promoter signal, have been modified by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence;

(3) a method for the production of recombinant influenza viruses as defined in (1) and (2) above comprising

- (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense, or in sense tandem design,
- (b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in the corresponding viral RNA segment, i.e., coding for the same viral gene as present in the tandem segment distal position, and
- (c) thereafter selective vRNA inactivation through ribozyme cleavage;

(4) a pharmaceutical composition comprising a recombinant influenza virus as defined in (1) and (2) above;

(5) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing a medicament for vaccination purposes;

(6) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for somatic gene therapy;

(7) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for transfer and expression of foreign genes into cells (abortively) infected by such viruses;

(8) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for transfer and expression of RNA molecules into cells infected by such viruses;

(9) a method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus as defined in (1) and (2) above as expression vector;

(10) a method for preventing and/or treating influenza which comprises administering a recombinant influenza virus as defined in (1) and (2) above to the mammal to be treated, i.e., a vaccination method utilizing said recombinant virus;

(11) a method for somatic gene therapy, which method comprises subjecting the organism to be treated with a recombinant influenza virus as defined in (1) and (2) above;

(12) a method for transfer and expression of foreign genes into cells, and for transfer and expression of RNA molecules into cells, which method comprises infecting the cells with a recombinant influenza virus as defined in (1) and (2) above;

(13) use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for autologous immunotherapy;

(14) a method for an immunotherapy which comprises *ex vivo* infection of immune cells with a recombinant influenza virus as defined in (1) and (2) above, and introduction of the transduced cells into the patient; and

(15) a method for the induction of antibodies which comprises utilizing a recombinant influenza virus as defined in (1) and (2) above as an immunogen.

[0011] The invention is described in more detail below.

Brief Description of the Figures

[0012] Fig. 1 shows the basepair substitution analysis according to the vRNA 'corkscrew' structure:

(A) 'Corkscrew' conformation of the vRNA promoter drawn against a schematic indication of interacting tripartite viral polymerase. Paired positions exchanged in individual experiments are indicated by numbers, nucleotides 3 or 8 are counted from the 3' end. pHL2024 containing promoter-up mutation '1104' is used as the reference construct (=100%) in all of the CAT assays, while pHL2428 represents the wild-type promoter structure.

(B) CAT analysis of a series of substitution variants in positions 3 and 8 from the 5' end as indicated above the lanes; 50 µl of cell lysate obtained from 10⁶ MDCK cells infected in the first viral passage with recombinant viral progeny.

(C) pHL2024 and pHL1920 comparative CAT analysis, in 100 fold dilution relative to (B), i.e., obtained from 0.5 µl of cell lysate in 3 h reaction time.

[0013] Fig. 2: Vector plasmid pHL1920, the exact sequence of the 3888 bps circular DNA is shown in SEQ ID NO: 20

[0014] Fig. 3 shows the genetic structure and the RNA transcription products of influenza model tandem expression constructs. Heavy lines for the plasmid cDNA constructs refer to double-stranded DNA, while single-stranded RNA molecules are represented by thin lines, and their 5' to 3' directionalities are marked by arrows. Standard modifications

at their 5' and 3' ends are indicated by a dot (5' cap structure) and A_n (3' polyadenylation), both are absent in the primary anti-sense transcription product, the viral RNA (vRNA), which is transcribed by cellular RNA polymerase I (RPol). Full-length mRNA, is synthesized by influenza viral polymerase (virPo), and a partial splice reaction results in a functional yield of shorter mRNA₂ molecules. While both of the reporter genes are indicated on the DNA level, together with the positions of splice donor (D) and acceptor (A) signal sequences as well as the promoter (p_i) and terminator (t_i) elements for RNA polymerase I start and stop, on the RNA level only those genes and splice signals are marked that are actually translated into protein or actively involved in splicing. The chloramphenicol-acetyltransferase gene (CAT) has been inserted in proximal position in pHL3196 and pHL3235, and in distal position in pHL3224 and pHL3236 (see Figs. 5 to 8), while green fluorescent protein (GFP) in each case is located in alternate location. All vRNA molecules - and hence, also the cDNA constructs - carry sequence variations at their 3' ends, which together constitute the 1104 promoter-up mutations: G3A, U5C, C8U (nucleotide positions counted from the 3' vRNA end). pHL3235 and pHL3236 vRNAs are extended in size by about 1000 nucleotides of untranslated sequence relative to pHL3196 and pHL3226: 2600 instead of 1600 nucleotides in lengths. For full-size representation of circular plasmid DNAs see Figs. 5-8, for CAT expression data of all infected by recombinant influenza viruses carrying the respective viral RNAs see Fig 4.

[0015] Fig. 4 shows the CAT assay results for the group of tandem vRNA plasmid constructs as described in the Example. In particular, the ratio between chloramphenicol (bottom line) and acetylchloramphenicol (upper three lines) in a flash-CAT assay, after the 2nd (A) and 4th (B) passage of recombinant viruses carrying the reportergene CAT, can be determined from said figure. The following constructs were utilized:

pHL1844 (control): monocistronic CAT-construct downstream of promoter variant 1104.

pHL3196: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 1530 nucleotides (not "extended"), see also Fig. 5.

pHL3235: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 2550 nucleotides ("extended"), see also Fig. 7.

pHL3224: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 1700 nucleotides (not "extended"), see also Fig. 6.

pHL3236: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 2720 nucleotides ("extended"), see also Fig. 8.

pHL2899: ambisense construct, p_v -CAT \rightarrow \leftarrow GFP- p_c resulting in an RNA having a total length of 1500 nucleotides.

pHL2960: ambisense construct, p_v -CAT \rightarrow \leftarrow GFP- p_c resulting in an RNA having a total length of 1500 nucleotides.

[0016] The five constructs on the left side were transfected into the cell DNA without the use of "booster" plasmides, the four constructs on the right side were, however, transfected with the "booster" plasmides, which gives a jump-start of the constructs due to recombinant vRNA amplification prior to helper virus injection, equivalent to an advantage of about two passages. The "booster" plasmides comprise expression constructs for the nucleoprotein as well as the three subunits of influenza viral polymerase, each downstream of an RNA polymerase II promoter and in an mRNA forming cassette.

While the ambisense construct having the CAT-reporter gene in the weaker position, i.e. behind the cRNA promoter (pHL2899), is only expressed moderately, this is not the case in the respective tandem construct having the CAT-reporter gene in the weaker position, viz. pHL3224 or pHL3236. Further, the "extension" of the vRNA by 1020 non-translated nucleotides (at the 3' end) is tolerated without significant decrease of expression (see pHL3235 versus pHL3196).

[0017] Fig. 5: Vector plasmid pHL3196, the exact sequence of the 4500 bps circular DNA is shown in SEQ ID NO:21.

[0018] Fig. 6: Vector plasmid pHL3224, the exact sequence of the 4721 bps circular DNA is shown in SEQ ID NO:22.

[0019] Fig. 7: Vector plasmid pHL3235, the exact sequence of the 5517 bps circular DNA is shown in SEQ ID NO:23.

[0020] Fig. 8: Vector plasmid pHL3236, the exact sequence of the 5699 bps circular DNA is shown in SEQ ID NO:24.

Detailed Description of the Invention

[0021] According to the present invention "influenza virus" embraces influenza A virus, influenza B virus and influenza C virus, with influenza A virus being preferred.

[0022] "Bicistronic" according to the present invention refers to a viral RNA segment, vRNA, cRNA or mRNA that includes two independent genes in covalent junction; in a preferred version one of these genes is of viral origin, while the other one codes for a foreign, recombinant gene product.

[0023] "Proximal" and "proximal position" according to the present invention refers to the 5' portion of one of the genes in the bicistronic viral mRNA, i.e., ahead (upstream) of the second gene in "distal position".

[0024] A "mammal" according to the present invention includes humans and animals. "Organism" embraces prokaryotic and eukaryotic systems as well as multicellular systems such as vertebrates (including mammals) and inverte-

brates.

[0025] "Infected cells" and "infecting cells" according to the present invention also include "abortively infected cells" and "abortively infecting cells", respectively.

[0026] In a preferred influenza virus according to embodiment (1) at least one of the regular viral RNA segments is replaced by a tandem RNA segment which contains one of the standard viral genes in distal position, and a foreign, recombinant gene in proximal position, both in anti-sense orientation, or vice-versa. It is moreover preferred that in the tandem RNA molecule said foreign recombinant gene is covalently bound to one of the viral genes while the original vRNA segment coding for the same gene is deleted from the recombinant virus by specific ribozyme cleavage.

[0027] The foreign gene(s) in tandem covalent junction with the viral gene(s) preferably code for proteins and/or glycoproteins which are secreted from cells infected with the recombinant virus, such as lymphokines, or code for glycoproteins that are incorporated into the virion as well as the plasma membrane of the infected cell. In another preferred embodiment the foreign gene(s) in tandem covalent junction with the viral gene(s) code for proteins or artificial polypeptides designed to support an efficient presentation of inherent epitopes at the surface of infected cells, for stimulation of B cell and/or T cell response. Such proteins or artificial polypeptides constitute for instance a tumor antigen or an artificial oligomeric series of T cell epitopes that have been identified within a polypeptide chain. Finally, the foreign gene(s) may be suitable for transfer and expression of RNA molecules, including antisense RNAs and ribozymes, into cells. Such recombinant influenza viruses are suitable for sequence specific gene silencing, for example by antisense or RNA interference mechanisms.

[0028] A preferred recombinant virus of the invention is where in the regular viral RNA segments one or both of the standard glycoproteins hemagglutinin and neuraminidase have been exchanged, preferably into fusion glycoproteins consisting of an anchor segment derived from hemagglutinin and an ectodomain obtained from the foreign source, viral or cellular, or in which such recombinant glycoprotein has been inserted as a third molecular species in addition to the remaining standard components.

[0029] As set forth in embodiment (2) above, a preferred recombinant virus of the invention is where the terminal viral RNA sequences, which are active as promoter signal, have been modified by nucleotide substitution in up to 5 positions, resulting in improved transcription rates (of both the vRNA promoter and in the cRNA promoter as present in the complementary sequence) as well as enhanced replication and/or expression rates relative to the wild-type sequence. Said modified terminal viral RNA sequences differ from the wild-type sequence in that in said tandem vRNA segment the 12 nucleotide conserved influenza 3' terminal sequence has been modified by replacement of one to three nucleotides occurring in said sequence at positions 3, 5 and 8 relative to the 3' end by other nucleotides provided that the nucleotides introduced in positions 3 and 8 are forming a base pair (i.e., if the nucleotide position 3 is G, than that in position 8 is C; if the nucleotide in position 3 is C, than that in position 8 is G; etc.).

[0030] The 3' conserved regions of the wild-type influenza virus have the following sequences:

Influenza A: (5')-CCUGCUUUUGCU-3'

Influenza B: (5')-NN(C/U)GCUUCUGCU-3'

Influenza C: (5')-CCUGCUUCUGCU-3'.

[0031] Moreover, the 13 nucleotide conserved influenza 5'-terminal sequence may be modified by replacement of one or two nucleotides occurring in said sequence as positions 3 and 8 by other nucleotides, again provided that the introduced nucleotides are forming a base pair. The 5' conserved regions of the wild-type influenza virus have the following sequences:

Influenza A: 5'-AGUAGAAACAAGG

Influenza B: 5'-AGUAG(A/U)AACA(A/G)NN

Influenza C: 5'-AGCAGUAGCAAG(G/A):

[0032] Preferred influenza viruses of the invention are those wherein in the 3' conserved region the replacements G3A and C8U have been performed, more preferred are those where also the replacement U5C has been performed (the above mutations are annotated relative to the 3' end; such counting from the 3' end is also indicated by a line on top of the digit, e.g., G $\bar{3}$ A). Another preferred influenza virus mutant comprises the 3'-terminal nucleotide sequence G3C, U5C and C8G (relative to the 3' end) resulting in the following 3' terminal nucleotide sequence (5')-CCUGGU-

UCUCCU-3'. Among the influenza viruses defined hereinbefore those having a 3'-terminal nucleotide sequence of (5') -CCUGUUUCUACU-3' are most preferred. In case of an influenza A virus the segment may further have the modifications U3A and A8U in its 5' terminal sequence, in case of influenza C it may have the modifications C3U and G8A in its 5' terminal sequence. The most preferred influenza viruses of the present invention comprise the following general structures:

Influenza A (mutant pHL1104):

5'-AGUAGAAACAAGGNNNU₅₋₆..(880-2300 ntds)..N'N'N'CCUGUUUCUACU-3'

Influenza A (mutant pHL1920):

5'-AGAAGAAUCAAGGNNNU₅₋₆..(880-2300 ntds)..N'N'N'CCUGUUUCUACU-3'

Influenza A (mutant pHL1948):

5'-AGUAGAAACAAGGNNNU₅₋₆..(880-2300 ntds)..N'N'N'CCUGGUUCUCCU-3'

Influenza B:

5'-AGUAG(A/U)AACA(A/G)NNNNNU₅₋₆..(880-2300 ntds)..N'N'N'N'N'(C/U)GUUUCUACU-3'

Influenza C:

5'-AGUAGUAAACAAG(G/A)GU₅₋₆..(880-2300 ntds)..CCCCUGUUUCUACU-3'

[0033] In the above structures the variables are defined as follows:

- (1) Underlined and enlarged letters show the required mutations relative to the wild-type sequence for preparing a promoter mutant with enhanced properties;
- (2) enlarged A in position 10 in the 5'-part of the sequence: unpaired A residue, bulge-forming;
- (3) (A/G) in one position: different isolates or single segments with variable sequence at the respective position, which are functionally interchangeable;
- (4) N and N': positions undefined, but base-paired relative to each other because of complementarity between the 5' and 3' termini, different among the 8 segments, but constant for each segment throughout all viral isolates;
- (5) (880-2300 ntds): the lengths of the viral RNA segments, in case of segments with foreign genes increased up to 4,000 nucleotides.

[0034] According to embodiments (1) to (3) the invention provides

- a stable recombinant influenza virus containing (up to) seven regular vRNA segments plus one (or more) additional bicistronic segment(s) coding for a foreign gene in covalent conjunction with one of the influenza genes, in tandem arrangement, and
- a method for the construction of stable recombinant influenza viruses through tandem arrangement of bicistronic vRNA segments, which is also applicable as a method for attenuation and for prevention of reassortment between co-infecting influenza viruses.

[0035] Expression of both gene products in these constructions is made possible by way of an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region of such a quality that splicing does occur in part of the mRNA molecules only, i.e., both mRNAs spliced and unspliced are present in the infected cell. For compensation with regard to the vRNA length the bicistronic segment is connected to a promoter variant of enhanced replication and transcription rates as defined herein before.

[0036] The splice donor and the splice acceptor signals are selected from authentic sequences as present in influenza segments 7 and 8 or other partially effective splice reaction substrates, preferably those of influenza virus WSN segment 7, i.e., 5'-AG↓GTACGTTTC-3' (donor) and 5'-GCTGAAAAATGATCTTCTTGAAAAATTGCAG↓GC-3' (acceptor).

[0037] In a particular application of embodiments (1) to (3) the tandem bicistronic mRNA codes for one of the viral genes, such as hemagglutinin, in conjunction with all or part of the viral neuraminidase coding sequence, in antisense orientation, while the authentic neuraminidase vRNA segment is missing in these recombinant viruses. In another variation of these constructs an anti-neuraminidase ribozyme sequence is also provided together with the (partial) neuraminidase antisense sequence, in the proximal position of these bicistronic recombinant segments. Recombinant viruses of this character are propagated in culture media with addition of exogenous neuraminidase.

[0038] The absence of a functional neuraminidase gene serves as a strong attenuation mechanism resulting in single-step infections of such recombinant viruses only. While a functional neuraminidase gene could be provided through another (wildtype) influenza virus superinfecting the same cell, expression of that gene is very much reduced through antisense RNA interaction and/or destruction of the corresponding vRNA through ribozyme cleavage, designed to interfere with production of infectious progeny even from co-infected cells; as a barrier against reassortment in double infected cells.

[0039] Recombinant viral RNAs coding simultaneously for two genes in tandem in a construct in which one of the viral genes is in covalent junction with a foreign coding sequence, are constructed via E. coli plasmid vector DNAs designed for an *in vivo* transcription of minus-strand vRNAs by cellular RNA polymerase I. In these constructs the gene in plus-strand proximal (upstream) position is surrounded by splice signals of limited activity such that both mRNAs, spliced and unspliced are present in the infected cell. Either the foreign gene or the viral gene may be in that upstream position. In the majority of applications the higher rates of expression will be reserved for the foreign coding sequence, while the lower expression rate of the viral gene is adapted to be approximately in balance with expression of the other viral genes encoded by the regular viral segments.

[0040] To achieve such a balanced rate of expression, the splice signals and the promoter have to be chosen properly (Flick and Hobom, Interaction of influenza virus, polymerase with viral RNA in the 'corkscrew' conformation, J. Gen. Virol. 80, 2565-2572 (1999)). At an increased overall transcription rate, the resulting mRNAs shall be spliced inefficiently if the viral gene is in the distal (downstream) position. Vice-versa, if the foreign gene is in the distal position, splicing to obtain the foreign mRNA shall be achieved efficiently. Both designs serve to reach an over-expression of the foreign gene relative to the viral gene, of which the expression shall be in balance with the expression of the other viral genes. Further, the promoter variant attached to the bicistronic segment has the function to compensate for the increased gene length by way of an increased replication rate.

[0041] The influenza vRNA segments preferably used for construction of bicistronic segments include the neuraminidase (No. 6), hemagglutinin (No. 4) and NS segment (No. 8). In the NS segment the foreign gene may also substitute for the NS1 gene leaving the viral NS2 gene in its place. These recombinant viruses can, as an example, be made by the following procedure: A recombinant virus population can be selected by repeated ribozyme-mediated cleavage of helper-virus segments carrying ribozyme cleavage sites that flank the same viral gene in the monocistronic segment as is present in the bicistronic construct (PCT/EP00/01903). By serial viral passaging and relying on the output of reporter genes in equivalently constructed bicistronic segments, a balanced mode of expression can be achieved in choosing the right set of elements: promoter, splice signals plus a limited variation in segment length. The construct that gives rise to the balanced, stable expression is then used as a basis for a multiple cDNA transfection procedure in a helper-virus free design according to Neumann et al., Proc. Natl. Acad. Sci. USA, Vol 96, 9345-9350 (August 1999). The resulting recombinant influenza virus, obtained via single plaques in pure helper-free state is subjected to another series of propagation steps to finally evaluate its properties.

[0042] In a particular application this design is used for a controlled mode of viral attenuation. Attenuation of influenza viruses so far has been achieved in cold-sensitive mutants (Edwards et al., J. Infect. Dis. 169, 68-76(1994)), by deletion of the NS1 gene (partial attenuation, Egorov et al., J. Virol. 72, 6437-6441 (August 1998) and Palese et al., Proc. Natl. Acad. Sci. USA, 4309-4314 (April 2000)), or through deletion of the neuraminidase gene (full attenuation, Kawaoka et al., J. Virol. 74, 5206-5212 (June 2000)). The latter approach is adapted here using a novel technique for the attenuation, which for the first time is also able to interfere with (chance) superinfection by wild-type viruses.

[0043] In this embodiment of the invention a bicistronic cDNA construct is achieved, which instead of a foreign gene is coding either for part of or for the entire viral neuraminidase gene in antisense orientation, with or without being surrounded both by splice donor and acceptor elements. In another version of that design a 2 x 50 nucleotide antisense segment complementary to the 5'-terminal neuraminidase sequence has been cloned in flanking positions relative to a ribozyme construct according to the hammerhead design and oriented against a common GUC triplet within the neuraminidase sequence. In a preferred design this antisense expression construct has been attached to the hemagglutinin vRNA segment, while another gene or reporter gene is encoded in a second bicistronic vRNA, in conjunction with NS2.

[0044] Propagation of recombinant viruses deleted for the neuraminidase (NA) gene requires an addition of external

neuraminidase to the medium. In the absence of neuraminidase, infection by the NA deletion viruses is abortive: no infectious progeny is produced. Upon co-infection (3:3) of recombinant viruses together with wildtype viruses no progeny virus or plaque is observed, which is attributed to antisense-blocked expression or (partial) destruction of the neuraminidase segment originating from the wild-type virus. Therefore, the recombinant viruses described are not only attenuated in single infections, but simultaneously interfere with wildtype virus superinfection, and therefore, no reassortment between the two viruses will occur.

[0045] The pharmaceutical composition according to embodiment (4) above and the medicament of embodiment (5) above contain the recombinant influenza virus in a pharmaceutically effective amount. Besides said recombinant influenza virus, the pharmaceutical composition and the medicament may contain further pharmaceutically acceptable carrier substances well-known to a person skilled in the art, such as binders, desintegrants, diluents, buffers, preservatives, etc. The pharmaceutical composition and medicaments are solid or liquid preparations and are suitable to be administered orally, intravenously or subcutaneously.

[0046] The medicament according to embodiment (5) above is preferably suitable as a medicament against influenza and/or against other infections. The recombinant influenza virus may be present in form of inactivated preparations or may be present in form of live recombinant viruses, preferably as attenuated viruses.

[0047] Live recombinant viral vaccines, live but attenuated recombinant viral vaccines or inactivated recombinant viral vaccine can be formulated. Inactivated vaccines are "dead" in the sense that their infectivity has been destroyed. Ideally, the infectivity is destroyed without affecting its immunogenicity. To prepare inactivated vaccines, the recombinant virus may be grown in cell cultures or in embryonated chicken eggs, purified, and inactivated by formaldehyde or β -propiolactone. The resulting vaccine is usually administered intramuscularly.

[0048] Inactivated viruses may be formulated with suitable adjuvants to enhance the immunological response. Such adjuvants include, but are not limited to, mineral gels, e.g., aluminum hydroxide, surface-active substances such as pluronic polyols, lysolecithin, peptides, oil emulsions, and potentially useful human adjuvants such as BCG.

[0049] Many methods may be used to introduce the vaccine formulations above, for example the oral, intradermal, intramuscular, intraperitoneal, subcutaneous, or intranasal routes. Where a live recombinant virus vaccine is used, it is preferred to introduce the formulation via the natural route of infection for influenza virus.

[0050] The medicament according to embodiment (5) above is preferably suitable for prophylactic or therapeutic vaccination, or both, against influenza and other infections. For example, recombinant viruses can be made for use in vaccines against HIV, hepatitis B virus, hepatitis C virus, herpes viruses, papilloma viruses, to name but a few. In one embodiment the recombinant virus contains the genes for surface proteins of the viruses, in another the genes for non-structural or regulatory genes. The recombinant viruses may be present in form of inactivated preparations or may be present in form of live recombinant viruses, or as live, but attenuated viruses. In an attenuated virus the recombinant virus would go through a single or at most very few propagation cycle(s) and induce a sufficient level of immune response, but would not cause disease. Such viruses lack one of the essential influenza genes or contain mutations to introduce temperature sensitivity.

[0051] The agents of embodiments (6)-(8) of the invention are applicable in *ex vivo* and *in vivo* application schemes. The RNA molecule to be expressed by means of the agent of the embodiment (8) is of an antisense sequence or double strand sequence (in ambisense bidirectional transcription) relative to a target cellular mRNA molecule. In embodiment (8) the agent is preferably suitable for sequence-specific gene silencing, preferably by antisense RNA or RNA interference mechanisms.

[0052] The method for the production of proteins or glycoproteins is preferably performed in cell culture cells or in fertilized chicken cells in accordance with standard techniques within the general knowledge of a person skilled in the art. The proteins or glycoproteins to be expressed are those incorporated into the ambisense construct as defined hereinbefore.

[0053] The methods according to embodiments (9) to (12), (14) and (15) of the invention include the administration of an effective amount to the mammal or the administration of a sufficient infective dose of the recombinant virus to the cell system that is used for *ex vivo* therapy or for *in vitro* investigations, whereby the amount and dose will be determined by a person skilled in the respective arts or knowledgeable of the desired treatments.

[0054] The agent of embodiment (14) of the invention is preferably utilized to infect, transfect or transduce patient-derived immune cells. The agent is suitable for treatment of cancer or chronic viral infections. For this purpose, patient derived immune cells, preferably dendritic cells, are *ex vivo* infected with recombinant influenza viruses expressing, e.g., tumor antigens or viral antigens. The transduced cells are then reintroduced into the patient.

[0055] The preferred method for immunotherapy of embodiment (14) of the invention is an autologous immunotherapy, wherein the cells which are *ex vivo* infected are patient-derived and the transduced cells are reintroduced into the patient. The diseases to be treated by this method include cancer and chronic viral infections. For details regarding such treatment see discussion of embodiment (13) above.

[0056] The method for inducing antibodies according to embodiment (15) of the invention is suitable for inducing antibodies to foreign proteins including glycoproteins, following the administration of protein or glycoprotein antigens

as part of a recombinant influenza virus in an authentic conformation, whereby the virus is purified by gentle procedures based on hemagglutination, and the gene is expressed at high rates in the infected cells.

[0057] As influenza viruses have a wide host range, recombinant influenza viruses can be used to obtain strong immune responses in, and isolate antibodies from, a wide range of animals, including, but not limited to, fowl, pigs, horses, and mice. Further, influenza viruses adapted to the mouse can be used for the infection of mice by several routes including the intranasal route. This results in infection of the pharyngeal mucosal cells and results in an additional type of B cell response (e.g., as recognized in the ratio of IgG to IgA). Mice are of particular utility in the induction of immune responses in transgenic mice that have been engineered to express human antibodies. As gentle procedures based on hemadsorption are used to purify influenza viruses, antibodies to antigens in native conformation can be isolated from the infected mammals.

The preset invention further illustrated by the following, non-limiting Example:

Example

[0058] Model tandem bicistronic expression constructs using reporter genes CAT and GFP.

Objective: Measurements of relative expression rates for CAT in proximal and distal position, with live observation of GFP fluorescence in alternate position during propagation of recombinant influenza viruses.

a) Construction of bicistronic expression plasmid DNAs:

[0059] Starting out with vector plasmid pHH10 (Hoffmann, Ph.D. Thesis, Univ. Giessen (1997)), i.e. an ampicillin resistant plasmid including in between a human rDNA promoter segment and a murine rDNA terminator segment precisely inserted cDNA sequence elements representing the 5' and 3' vRNA sequence of influenza rRNA segment 5, and finally a central multiple cloning site sequence as obtained from plasmid PBSK, both reporter genes have been inserted in a stepwise manner. After that, to the proximal reading frame, i.e. CAT in pHL3196, and GFP in pHL3224, has been added an upstream splice donor sequence element and a downstream splice acceptor element, both inserted as double-strand oligonucleotides, in between particular restriction cleavage sites available in the respective positions. The signal sequences used in that pair of plasmids indicated above have been derived from influenza vRNA segment 7, which is known for its partial splice reactions yielding both gene products, M1 and M2, simultaneously. By insertion of a non-transcribed DNA fragment (representing an internal segment of the influenza PB1 coding region) in a distal position relative to both reading frames, pHL3196 has been converted into pHL3235, and pHL3224 into pHL3236. For the resulting plasmid constructs see Figs 5-8 and SEQ ID NOs: 21-24.

b) Transfection of plasmid DNAs and isolation of recombinant influenza viruses:

[0060] Semi-confluent 293-T cells, a human renal carcinoma cell line carrying an artificially integrated tumor virus SV40 T-antigen gene, were DNA-transfected using lipofectamine: 5-10 µg of DNA mixed with 30 µl Lipofectamine® (GIBCO/BRL) were added to 370 µl of DMEM medium and were incubated with 5x10⁶ to 10⁷ cells, washed and maintained serum-free for 5 to 8 hours, before serum was added for another 12 to 15 hours. Finally influenza helper virus FPV_{Bratislava} was used for infection of the DNA-transfected cells. The supernatant containing a mixture of helper viruses and recombinant viruses was collected for further propagation after 8 to 12 hours of infection, while the sedimented cells were used for preparation of a cell lysate, fractions of which were inserted in the CAT assay procedure. Viral propagation was achieved by infection of MDCK cells (Madin-Darby canine kidney cell line) again in semi-confluent state (5x10⁶ to 10⁷ cells per plate), generally using 1ml of the previous supernatant for infection. Serial propagations were done in the same way, with preparation of cell lysates for CAT assays at the end of each step. Infected cells were also used for observation of GFP fluorescence.

c) CAT assay:

[0061] Bacterial chloramphenicol-acetyltransferase (CAT) is accumulated in eukaryotic cells without degradation and can be used for representative gene expression measurements. The substrate used here is fluorescent boron-dipyrromethane-chloramphenicol difluoride (FLASH CAT-KIT®; Stratagene). 50 µl of cell lysate or reduced/diluted samples thereof were used for incubation with 7.5 µl of fluorescent substrate and 10 µl acetyl-CoA (4mM) co-substrate in 19 mM Tris/HCl, pH: 7.5 at 37°C for 3 hours. For extraction of reaction products 1ml of ethylacetate ins added, the mixture is vortexed, and separated by centrifugation. After solvent evaporation and dissolution again in 20ml ethylacetate, the reaction products are separated on a silica thin-layer chromatography plate using chloroform/methanol 87:13% (vol.) and the results are documented by photography under UV light.

d) Results

[0062] CAT in proximal or in distal position of this pair of recombinant plasmids is expressed about equally (Fig.4), and the same is true for GFP (not shown). The expression rates are increasing during the initial steps of viral propagation and stay about constant afterwards during further steps of recombinant viral passages, different from expression rates in ambisense bicistronic constructs (pHL2899 and pHL2960) (Fig. 4B). Co-transfection of booster plasmids in the initial 293-T cells increase the yields of recombinant viruses within the progeny population, which are maintained during consecutive steps of propagation. Addition of 1000 nucleotides of untranslated vRNA sequence will not reduce the expression rates substantially (pHL3235 versus pHL3196, and pHL3236 versus pHL3224).

EP 1 174 514 A1

SEQUENCE LISTING

<110> ARTEMIS PHARMACEUTICALS GmbH

5 <120> Recombinant Influenza Viruses with Bicistronic vRNAs Coding for
Two Genes in Tandem Arrangement

<130> 000746ep/JH/ml

10 <140>
<141>

<160> 24

15 <170> PatentIn Ver. 2.1

<210> 1
<211> 12
20 <212> RNA
<213> Influenza A virus

<400> 1
25 ccugcuuuug cu 12

<210> 2
<211> 12
30 <212> RNA
<213> Influenza B virus

<400> 2
35 nnygcuucug cu 12

<210> 3
<211> 12
40 <212> RNA
<213> Influenza C virus

<400> 3
45 ccugcuucug cu 12

<210> 4
<211> 12
<212> RNA
50 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Modified
influenza A 5'-sequence (pHL1104 and pHL1920)

55

EP 1 174 514 A1

	<400> 4	
	ccuguuucua cu	12
5	<210> 5	
	<211> 12	
	<212> RNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Description of Artificial Sequence: Modified	
	influenza A 3'-sequence (pHL1948)	
15	<400> 5	
	ccugguucuc cu	12
20	<210> 6	
	<211> 13	
	<212> RNA	
	<213> Influenza A virus	
25	<220>	
	<223> Description of Artificial Sequence: Modified	
	influenza A 3'-sequence (pHL1948)	
30	<400> 6	
	aguagaaaca agg	13
35	<210> 7	
	<211> 13	
	<212> RNA	
	<213> Influenza B virus	
40	<400> 7	
	aguagwaaca rnn	13
45	<210> 8	
	<211> 13	
	<212> RNA	
	<213> Influenza C virus	
50	<400> 8	
	agcaguagca agr	13
55	<210> 9	
	<211> 13	
	<212> RNA	
	<213> Artificial Sequence	

EP 1 174 514 A1

<220>
 <223> Description of Artificial Sequence: Modified
 influenza A 5'-sequence (pHL1920)
 5
 <400> 9
 agaagaauca agg 13
 10
 <210> 10
 <211> 21
 <212> RNA
 <213> Influenza A virus
 15
 <400> 10
 aguagaaaca aggnnnuuuu u 21
 20
 <210> 11
 <211> 21
 <212> RNA
 <213> Artificial Sequence
 25
 <220>
 <223> Description of Artificial Sequence: Modified
 influenza A 5'-sequence (pHL1920)
 30
 <400> 11
 agaagaauca aggnnnuuuu u 21
 35
 <210> 12
 <211> 21
 <212> RNA
 <213> Influenza B virus
 40
 <400> 12
 aguagwaaca rnnnnuuuu u 21
 45
 <210> 13
 <211> 19
 <212> RNA
 <213> Artificial Sequence
 50
 <220>
 <223> Description of Artificial Sequence: Modified
 influenza C 5'-sequence
 <400> 13
 aguaguaaca agrguuuuu 19
 55

EP 1 174 514 A1

5 <210> 14
 <211> 15
 <212> RNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Modified
 influenza A 3'-sequence (pHL1104 and pHL1920)
 10
 <400> 14
 nnnccuguuu cuacu 15

 15 <210> 15
 <211> 15
 <212> RNA
 <213> Artificial Sequence
 20
 <220>
 <223> Description of Artificial Sequence: Modified
 influenza A 3'-sequence (pHL1948)
 25 <400> 15
 nnnccugguu cuccu 15

 30 <210> 16
 <211> 15
 <212> RNA
 <213> Artificial Sequence
 35 <220>
 <223> Description of Artificial Sequence: Modified
 influenza B 3' sequence
 <400> 16
 40 nnnnnyguuu cuacu 15

 45 <210> 17
 <211> 14
 <212> RNA
 <213> Artificial Sequence
 50 <220>
 <223> Description of Artificial Sequence: Modified
 influenza C 3'-sequence
 <400> 17
 ccccguguuc uacu 14
 55

EP 1 174 514 A1

<210> 18
 <211> 10
 <212> DNA
 5 <213> Influenza A virus

 <400> 18
 aggtacggttc 10

 10
 <210> 19
 <211> 32
 <212> DNA
 <213> Influenza A virus

 15
 <400> 19
 gctgaaaaat gatcttcttg aaaattgcag gc 32

 20
 <210> 20
 <211> 3888
 <212> DNA
 <213> Artificial Sequence

 25
 <220>
 <223> Description of Artificial Sequence: PHL1920

 <400> 20
 30 cccaaaaaaa aaaaaaaaaa aaaaaaaaaa agtccagagt ggccccgccc ttccgcgccc 60
 gggggggggg ggggggggga cacttttcgga catctggtcg acctccagca tcgggggaaa 120
 aaaaaaaaac aaagttttcgc ccggagtagt ggtcgacctc cgaagtggg ggggagtaga 180
 aacagggttag ataatacactc actgagtgac atccacatcg cgagcgcgcg taatacgact 240
 cactatagggt cgaattgggt accgggcccc cctcagagt cgacggtatc gataagcttc 300
 gacgagattt tcaggagcta aggaagctaa aatggagaaa aaaatcactg gatataccac 360
 35 cgttgatata tcccaatggc atcgtaaaga acattttgag gcatttcagt cagttgctca 420
 atgtacctat aaccagaccg ttcagctgga tattacggcc tttttaaaga ccgtaaagaa 480
 aaataagcac aagtttttatc cggcctttat tcacattctt gcccgccctga tgaatgctca 540
 tccggaattc cgtatggcaa tgaaagacgg tgagctgggt atatgggata gtgttcaccc 600
 ttgttacacc gttttccatg agcaaaactga aacgttttca tcgctctgga gtgaatacca 660
 40 cgacgatttc cggcagtttc tacacatata ttcgcaagat gtggcgtgtt acggtgaaaa 720
 cctggcctat ttccctaag ggttttattga gaatatgttt ttctgtctcag ccaatccctg 780
 ggtgagtttc accagttttg atttaaactg ggccaatatg gacaacttct tcgccccctg 840
 tttcaccatg ggcaaatatt atacgcaagg cgacaagggt ctgatgccgc tggcgattca 900
 45 ggttcacatc gccgtttgtg atggcttcca tgcggcaga atgcttaatg aattacaaca 960
 gtactgcgat gagtggcagg gcggggcgta atttttttaa ggcagttatt ggtgccctta 1020
 aacgcctggt gctacgcctg aataagtgat aataagcggg tgaatggcag aaattcgctg 1080
 aagcttgata tcgaattcct gcagcccggg ggatccacta gttctagagc ggccgccacc 1140
 gcggtgggagc tcagcttttt gtccctttta gtgagggtta attgcgcgca ggcctagcta 1200
 50 ggtaaagaaa aataacccttg attcttctaa taaccggcg gcccaaatg ccgactcgga 1260
 gcgaaagata tacctcccc ggggccggga ggtcgcgtca ccgaccagc cgcgggcccc 1320
 ggcgacgcgc gacacggaca cctgtcccca aaaacgccac catcgagcc acacacggag 1380
 cggccggggc cctctggtca accccaggac acacgcggga gcagcgcgg gcccgggacg 1440
 cctcccggc cgcctgtgcc acacgcaggg ggccggcccc tgtctccaga gcgggagccg 1500
 55 gaagcatttt cggccggccc ctctacgac cgggacacac gagggaccga agcccggcc 1560

EP 1 174 514 A1

```

5  ggcgcgacct ctccgggccc acgcgcgcctc agggagcgct ctccgaactcc gcacggggac 1620
   tcgccagaaa ggatcgtgac ctgcattaat gaatcagggg ataacgcagg aaagaacatg 1680
   tgagcaaaaag gccagcaaaa ggccagggaac cgtaaaaagg ccgcgttgct ggcgtttttc 1740
   cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 1800
   aacccgacag gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct 1860
   cctgttccga cctgcccgt taccggatac ctgtccgctt ttctcccttc gggaagcgtg 1920
   gcgctttctc atagctcacg ctgtaggtat ctgagttcgg ttaggtcgt tcgctccaag 1980
   ctgggctgtg tgcacgaacc ccccgctcag cccgaccgct gcgccttacc cggtaactat 2040
10  cgtcttgagt ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac 2100
   aggattagca gagcgaggta ttaggcgggt gctacagagt tcttgaagtg gtggcctaac 2160
   tacggctaca ctagaaggac agtatttggg atctgcgctc tgctgaagcc agttaccttc 2220
   ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggttt 2280
   tttgtttgca agcagcagat tacgcgcaga aaaaaggat ctcaagaaga tcttttgatc 2340
15  ttttctacgg ggtctgacgc tcagtggaac gaaaactcac gttaagggat tttggtcatg 2400
   agattatcaa aaaggatctt cacctagatc cttttaaat aaaaatgaag ttttaaatca 2460
   atctaaagta tatatgagta aacttgggtc gacagttacc aatgcttaat cagtgaggca 2520
   cctatctcag cgatctgtct atttcgttca tccatagttg cctgactccc cgtcgtgtag 2580
   ataactacga tacgggaggg ttaccatct ggcccagtg ctgcaatgat accgcgagac 2640
20  ccacgctcac cgggtcccga tttatcagca ataaaccagc cagccggaag ggccgagcgc 2700
   agaagtggtc ctgcaacttt atccgcctcc atccagtcta ttaattgttg ccgggaagct 2760
   agagtaagta gttcgccagt taatagtttg cgcaacgttg ttgccattgc tacaggcatc 2820
   gtggtgtcac gctcgtcgtt tggtaggct tcattcagct ccggttccca acgatcaagg 2880
   cgagttacat gatcccccac gttgtgcaaa aaagcgggta gtccttcggt tctccgac 2940
25  gttgtcagaa gtaagttggc cgcagtgtta tcaactcatg ttatggcagc actgcataat 3000
   tctcttactg tcatgccatc cgtaagatgc tttctgtgta ctggtgagta ctcaaccaag 3060
   tcattctgag aatagtgtat gcggcgaccg agttgctctt gcccggcgct aacacgggat 3120
   aataccgcgc cacatagcag aactttaaaa gtgctcatca ttggaaaacg ttcttcgggg 3180
   cgaaaactct caaggatctt accgctgttg agatccagtt cgatgtaacc cactcgtgca 3240
30  cccaactgat cttcagcatc ttttactttc accagcggtt ctgggtgagc aaaaacagga 3300
   aggcaaaatg ccgcaaaaaa ggggaataagg gcgacacgga aatggtgaat actcactc 3360
   ttcttttttc aatattattg aagcatttat cagggttatt gtctcatgag cggatacata 3420
   tttgaatgta tttagaaaaa taaacaaaag agttttaga aacgcaaaaa ggccatccgt 3480
   caggatggcc ttctgcttaa tttgatgcct ggaggtttat ggccggcgct ctgcccgcga 3540
35  cctccggggc cgttgcttcg caacgttcaa atccgctccc ggccgatttg tctactcag 3600
   gagagcgttc accgacaaac aacagataaa acgaaaggcc cagtctttcg actgagcctt 3660
   tcggtttatt tgatgcctgg cagttcccta ctctcgcatg gggagacccc acactaccat 3720
   cggcgctacg gcgtttcact tctgagttcg gcatggggtc aggtgggacc accgcgtac 3780
   tgccgccagg caaattctgt tttatcagac cgcttctgct ttctgattta atctgtatca 3840
40  ggctgaaaat cttctctcat ccgcaaaaac agaagctagc ggccgac 3888

```

<210> 21

<211> 4500

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: pHL3196

<400> 21

```

agtagaaaca gggtagataa tcaactcactg agtgacatcc acatcgcgag cgcgaaggta 60
cgttctcgag cgcgcgtaat acgactcact ataggcgcaa ttgggtacgt tccatcatgg 120
agaaaaaaat cactggatat accaccgttg atatatccca atggcatcgt aaagaacatt 180

```

EP 1 174 514 A1

	ttgaggcatt	tcagtcagtt	gctcaatgta	cctataacca	gaccgttcag	ctggatatta	240
	cggccttttt	aaagaccgta	aagaaaaata	agcacaagtt	ttatccggcc	tttattcaca	300
	ttcttgcccg	cctgatgaat	gctcatccgg	aattccgtat	ggcaatgaaa	gacggtgagc	360
5	tgggtgatatg	ggatagtgtt	cacccttggt	acaccgtttt	ccatgagcaa	actgaaacgt	420
	tttcatcgct	ctggagtga	taccacgacg	atctccggca	gtttctacac	atatattcgc	480
	aagatgtggc	gtgttacgg	gaaaacctgg	cctatttccc	taaagggttt	attgagaata	540
	tgtttttcgt	ctcagccaat	ccctgggtga	gtttcaccag	ttttgattta	aacgtggcca	600
	atatggacaa	cttcttcgcc	cccgttttca	ccatgggcaa	atattatacg	caaggcgaca	660
10	aggtgctgat	gccgctggcg	attcaggttc	atcatgccgt	ctgtgatggc	ttccatgtcg	720
	gcagaatgct	taatgaatta	caacagtact	gcgatgagtg	gcagggcggg	gcgcgttaac	780
	gagatcagct	gaaaaatgat	cttcttgaaa	atttgcaggc	cgtacgtgta	ccgggcccc	840
	cctcgactcg	cgaaggagtc	caccatgagt	aaaggagaag	aacttttcac	tggagttgtc	900
	ccaattcttg	ttgaattaga	tgggtgatgtt	aatgggcaca	aattttctgt	cagtggagag	960
15	ggtgaagggtg	atgcaacata	cggaaaactt	acccttaa	ttatttgcac	tactggaaaa	1020
	ctacctgttc	catggccaac	acttgtcact	actttcactt	atgggtgttca	atgcttttca	1080
	agatacccag	atcatatgaa	acagcatgac	tttttcaaga	gtgccatgcc	cgaagggtat	1140
	gtacaggaaa	gaactatatt	tttcaaatg	gacgggaact	acaagacacg	tgctgaagtc	1200
	aagtttgaag	tgatatacc	tgttaataga	atcgagttaa	aagggtattga	ttttaaagaa	1260
20	gatggaaaaca	ttcttgga	caaattggaa	tacaactata	actcacacaa	tgtatacatc	1320
	atggctgaca	agcagaagaa	cggaatcaag	gccaacttca	agaccgcga	caacatcgag	1380
	gacggcgggc	tgcagctggc	cgaccactac	cagcagaaca	ccccaattgg	cgatggccct	1440
	gtcctttttac	cagacaacca	ttaqctgtcc	acacaatctg	ccctttcgaa	agatcccaac	1500
	gaaaagagag	accacatggt	ccttcttgag	tttgtaacag	ctgctgggat	tacacatggc	1560
25	atggatgaac	tatacaagg	atcccatcac	catcaccatc	actaagctcc	atgggtctaga	1620
	tatcgatagg	cctagctagg	taaagaaaa	taccttgtt	tctactaata	acccggcggc	1680
	ccaaaatgcc	gactcggagc	gaaagatata	cctccccggg	ggccgggagg	tcgcgtcacc	1740
	gaccacgcgc	ccggcccagg	cgacgcgcga	cacggacacc	tgtcccaaaa	aacgccacca	1800
	tcgcagccac	acacggagcg	cccggggccc	tctggtcaac	cccaggacac	acgcgggagc	1860
30	agcgccgggc	cggggacgcc	ctcccgcccg	cccgtgccac	acgcaggggg	ccggcccgtg	1920
	tctccagagc	gggagccgga	agcattttcg	gccggccctt	cctacgaccg	ggacacacga	1980
	gggaccgaag	gccggccagg	cgcgacctct	cgggcccgcac	gcgcgctcag	ggagcgctct	2040
	ccgactccgc	acggggactc	gccagaaagg	atcgtgacct	gcattaatga	atcaggggat	2100
	aacgcaggaa	agaacatgtg	agcaaaaagg	cagcaaaaagg	ccaggaaccg	taaaaaggcc	2160
35	gcgttgctgg	cgtttttcca	taggtccgc	cccctgacg	agcatcaca	aaatcgacgc	2220
	tcaagtcaga	ggtggcgaaa	cccagacgga	ctataaagat	accaggcggt	tccccctgga	2280
	agtcctctcg	tgcgctctcc	tgttccgacc	ctgccgctta	ccggataacct	gtccgccttt	2340
	ctcccttcgg	gaagcgtggc	gctttctcat	agctcacgct	gtagggtatct	cagttcgggtg	2400
	taggtcgttc	gtcccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	2460
40	gccttatccg	gtaactatcg	tcttgagtcc	aaccgggtaa	gacacgactt	atcgccactg	2520
	gcagcagcca	ctggtaacag	gattagcaga	gcgaggtatg	taggcgggtg	tacagagttc	2580
	ttgaagtgg	ggcctaacta	cggctacact	agaaggacag	tatttggtat	ctgcgctctg	2640
	ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	2700
	gctggtagcg	gtggtttttt	tgtttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	2760
45	caagaagatc	ctttgatctt	ttctacgggg	ctgacgctc	agtggaaacga	aaactcacgt	2820
	taagggattt	tggctcatgag	attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	2880
	aaatgaagtt	ttaaatcaat	ctaaagtata	tatgagtaaa	cttgggtctga	cagttacca	2940
	tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcac	catagttgcc	3000
	tgactccccg	tctgttagat	aaactacgata	cgggagggtt	taccatctgg	ccccagtgtc	3060
50	gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	aaaccagcca	3120
	gccggaaggg	ccgagcgag	aagtggtcct	gcaactttat	ccgcctccat	ccagtctatt	3180
	aattggttgc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	caacgttgtt	3240
	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc	3300
55	ggttcccaac	gatcaaggcg	agttacatga	tcccccatgt	tgtgcaaaaa	agcgggttagc	3360

EP 1 174 514 A1

5 tccttcggtc ctccgacgt tgtcagaagt aagttggcgg cagtgttatc actcatggtt 3420
 atggcagcac tgcataattc tcttactgtc atgccatccg taagatgctt ttctgtgact 3480
 ggtgagtact caaccaagtc attctgagaa tagtgtatgc ggcgaccgag ttgctcttgc 3540
 ccggcggtcaa cacgggataa taccgcgcca catagcagaa ctttaaaagt gctcatcatt 3600
 ggaaaaacgtt ctccggggcg aaaactctca aggatcttac cgctgttag atccagttcg 3660
 atgtaaccca ctctgtgcacc caactgatct tcagcatctt ttactttcac cagcgtttct 3720
 ggggtgagcaa aaacaggaag gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa 3780
 tgttgaatac tcatactctt cctttttcaa tattattgaa gcatttatca gggttatctg 3840
 10 ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaaagag tttgtagaaa 3900
 cgcaaaaagg ccatcctgca ggtatggcctt ctgcttaatt tgatgcctgg cagtttatgg 3960
 cgggcgtect gcccgccacc ctccgggcgg ttgcttcgca acgttcaa ccgtcccgg 4020
 cggatttgct ctactcagga gagcgttcac cgacaaacaa cagataaaac gaaaggccca 4080
 gtctttcgac tgagcctttc gttttatttg atgcctggca gtccctact ctcgcatggg 4140
 15 gagacccac actaccatcg ggcgtacggc gtttacttct tgagttcggc atggggtcag 4200
 gtgggaccac cgcgtactg ccgccaggca aattctgttt tatcagaccg cttctcgctt 4260
 ctgatttaatt ctgtatcagg ctgaaaaatc tctctcatcc gccaaaacag aagctagcgg 4320
 ccgatcccca aaaaaaaaaa aaaaaaagtc cagagtggcc ccgccgttcc 4380
 gccgcggggg gggggggggg gggggacact ttcggacatc tggtcgacct ccagcatcgg 4440
 20 gggaaaaaaa aaaaacaaag tttcgcccg agtactggtc gacctccgaa gttggggggg 4500

25 <210> 22
 <211> 4721
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> Description of Artificial Sequence: pHL3224

35 <400> 22
 atctagacca tggagcttag tgatggtgat ggtgatggga tcccttgat agttcatcca 60
 tgccatgtgt aatcccagca gctgttacaa actcaagaag gaccatgtgg tctctctttt 120
 cgttgggatac ttctgaaagg gcagattgtg tggacaggta atggttgctt ggtaaaaggga 180
 caggggccatc gccaatggg gtgttctgct ggtagtggtc ggccagctgc acgccgccgt 240
 40 cctcgatgtt gtggcggttc ttgaagtgtg ccttgatttc gttctcttgc ttgtcagcca 300
 tgatgtatac atttgtgtgag ttatagttgt attccaattt gtgtccaaga atgtttccat 360
 cttctttaaa atcaatacct ttttaactga ttctattaac aagggtatca ccttcaaact 420
 tgacttcagc acgtgtcttg tagttcccgt catctttgaa aaatatagtt ctttctctga 480
 cataaccttc gggcatggca ctcttgaaaa agtcatgctg tttcatatga tctgggtatc 540
 ttgaaaagca ttgaacacca taagtgaag tagtgacaag tgttggccat ggaacaggta 600
 gttttccagt agtgcaaaata aatttaaggg taagttttcc gtatgttgca tcaccttcac 660
 cctctccact gacagaaaat ttgtgcccac taacatcacc atctaattca acaagaattg 720
 ggacaactcc agtgaagagt tcttctcttc tactcatggt ggactccttc gcgagtcgag 780
 45 gggggggccc gtacacgtac gcgctcgaga acgtaccttc gcgctcgaga tgtggatgtc 840
 actcagttag tgattatcta cctgttttct actccccc aacttcggag gtcgaccagt 900
 actccgggcy aaactttgtt tttttttttt cccccgatgc tggaggtcga ccagatgtcc 960
 gaaagtgtcc cccccccccc ccccccccg cgcggaaagg cggggccact ctggactctt 1020
 tttttttttt tttttttttt ttttggggat cgcccgctag cttctgtttt ggcggatgag 1080
 50 agaagatttt cagcctgata cagattaaat cagaacgcag aagcggctct ataaaacaga 1140
 atttgctgtg cggcagtagc gcggtggctc cacctgaccc catgccgaac tcagaagtga 1200
 aacgccgtag cgcgatggg agtgtggggg ctccccatgc gagagtaggg aactgccagg 1260
 catcaaataa aacgaaaggc tcagtcgaaa gactgggctt ttcgttttat ctgttggttg 1320
 tcggtgaacg ctctcctgag taggacaaat ccgccgggag cggatttgaa cgttgcgagg 1380

EP 1 174 514 A1

	caacggcccc	gaggggtggcg	ggcaggacgc	ccgccataaa	ctgccaggca	tcaaattaag	1440
	cagaaggcca	tcctgacgga	tggccttttt	gcgtttctac	aaactctttt	gtttattttt	1500
	ctaaatacat	tcaaatatgt	atccgctcat	gagacaataa	ccctgataaa	tgcttcaata	1560
5	atattgaaaa	aggaagagta	tgagtattca	acatttccgt	gtcgccctta	ttcccttttt	1620
	tgccggcattt	tgcccttcctg	tttttgctca	cccagaaaacg	ctggtgaaag	taaaagatgc	1680
	tgaagatcag	ttgggtgcac	gagtggttta	catcgaaactg	gatctcaaca	gcggttaagat	1740
	ccttgagagt	tttcgccccg	aagaacgttt	tccaatgatg	agcactttta	aagtctctgt	1800
	atgtggcgcg	gtattatccc	gtgttgacgc	cgggcaagag	caactcggtc	gccgcataca	1860
10	ctattctcag	aatgacttgg	ttgagtactc	accagtcaca	gaaaagcatc	ttacggatgg	1920
	catgacagta	agagaattat	gcagtgcctg	cataaccatg	agtataaca	ctgcggccaa	1980
	cttacttctg	acaacgatcg	gaggaccgaa	ggagctaacc	gcttttttgc	acaacatggg	2040
	ggatcatgta	actcgccctg	atcggtggga	accggagctg	aatgaagcca	taccaaacga	2100
	cgagcgtgac	accacgatgc	ctgtagcaat	ggcaacaacg	ttgcgcaaac	tattaactgg	2160
15	cgaactactt	actctagctt	cccggcaaca	attaatagac	tggatggagg	cggataaagt	2220
	tgccaggacca	cttctgcgct	cggcccttcc	ggctggctgg	tttattgctg	ataaatctgg	2280
	agccgggtgag	cgtgggtctc	gcgggtatcat	tgccagcactg	gggccagatg	gtaagccctc	2340
	ccgtatcgta	gttatctaca	cgacggggag	tcaggcaact	atggatgaac	gaaatagaca	2400
	gatcgctgag	ataggtgcct	cactgattaa	gcattggtaa	ctgtcagacc	aagtttactc	2460
20	atatatactt	tagattgatt	taaaacttca	tttttaattt	aaaaggatct	aggtgaagat	2520
	cttttttgat	aatctcatga	ccaaaatccc	ttaacgtgag	ttttcgttcc	actgagcgtc	2580
	agaccccgta	gaaaagatca	aaggatcttc	ttgagatcct	tttttctgc	gcgtaatctg	2640
	ctgcttgcaa	acaaaaaac	caccgctacc	agcgggtggt	tgtttgccgg	atcaagagct	2700
	accaactctt	tttccgaagg	taactggctt	cagcagagcg	cagataccaa	atactgtcct	2760
25	tctagtgtag	ccgtagttag	gccaccactt	caagaactct	gtagcaccgc	ctacatacct	2820
	cgctctgcta	atcctgttac	cagtggctgc	tgccagtggc	gataagtcgt	gtcttaccgg	2880
	gttggtactca	agacgatagt	taccggataa	ggcgcagcgg	tcgggctgaa	cgggggggtt	2940
	gtgcacacag	cccagcttgg	agcgaacgac	ctacaccgaa	ctgagatacc	tacagcgtga	3000
30	gctatgagaa	agcgcacacg	ttcccgaagg	gagaaaaggcg	gacaggatc	cggtaagcgg	3060
	cagggtcgga	acaggagagc	gcacgaggga	gcttccaggg	ggaaacgcct	ggtatcttta	3120
	tagtcctgtc	gggtttcgcc	acctctgact	tgagcgtcga	tttttgat	gctcgtcagg	3180
	ggggcgaggc	ctatggaaaa	acgccagcaa	cgcggccttt	ttacggttcc	tggecttttg	3240
	ctggcctttt	gctcacatgt	tctttctgc	gttatccct	gattcattaa	tgccaggtcac	3300
35	gatcctttct	ggcgagtccc	cgtgaggagt	cggagagcgc	tccttgagcg	cgcgtgcggc	3360
	ccgagaggtc	gcgcctggcc	ggccttcggt	ccctcgtgtg	tcctcgctgt	aggaggggcc	3420
	ggccgaaaat	gcttccggct	cccgtctctg	agacacgggc	cggcccccctg	cgtgtggcac	3480
	ggggcgccgg	gagggcgctc	cgggcccggc	gctgtctccc	cgtgtgtcct	gggggtgacc	3540
	agagggcccc	gggcgctccg	tgtgtggctg	cgatggtggc	gtttttgggg	acagggtgtc	3600
40	gtgtcgcgcg	tcgcctgggc	cggcggcgtg	gtcggtgacg	cgacctccc	gccccggggg	3660
	aggtatatct	ttcgctccga	gtcggcattt	tgggcccgcg	ggttattagt	agaaacaagg	3720
	gtatttttct	ttacctagct	aggcctgcgc	gcaattaacc	ctcactaaa	ggaacaaaag	3780
	ctggagctcc	accgcggtgg	cggccgctct	agaactagt	gatcccccg	gctgcaggaa	3840
	ttcgatatca	agcttcgacg	aatttctgcc	attcatccgc	ttattatcac	ttattcaggc	3900
45	gtagcaccag	gcgtttaagg	gcaccaataa	ctgccttaaa	aaaattacgc	cccgccttgc	3960
	cactcatcgc	agtactgttg	taattcatta	agcattctgc	cgacatggaa	gccatcacia	4020
	acggcatgat	gaacctgaat	cgcagcggc	atcagcacct	tgctgccttg	cgtataatat	4080
	ttgcccatgg	tgaaaacggg	ggcgaagaag	ttgtccatat	tggccacgtt	taaatcaaaa	4140
	ctggtgaaac	tcaccagggg	attggtctgag	acgaaaaaca	tattctcaat	aaaccttcta	4200
50	gggaaatagg	ccaggttttc	accgtaacac	gccacatctt	gcgaatatat	gtgtagaaac	4260
	tgccggaat	cgtcgtggta	ttcactccag	agcgatgaaa	acgtttcagt	ttgctcatgg	4320
	aaaacggtgt	aacaagggtg	aacactatcc	catatcacca	gctcaccgtc	tttcattgcc	4380
	atacgggaatt	ccggatgagc	attcatcagg	cgggcaagaa	tgtgaataaa	ggccggataa	4440
	aacttgtgct	tatttttctt	tacggtcttt	aaaaaggccg	taatatccag	ctgaacggtc	4500
55	tggttatagg	tacattgagc	aactgactga	aatgcctcaa	aatgttcttt	acgatgccat	4560

EP 1 174 514 A1

tgggatatat caacggtggt atatccagtg atttttttct ccatttttagc ttccttagct 4620
cctgaaaatc tcgtcgaagc ttatcgatac cgtcgacctc gagggggggc ccggtacggc 4680
ctgcaaattt tcaagaagat cattttttcag ctgatctcgt t 4721

5

<210> 23

<211> 5517

<212> DNA

10

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: pHL3235

15

<400> 23

20

25

30

35

40

45

50

55

agtagaaaaca gggtagataa tcactcactg agtgacatcc acatcgcgag cgcgaaggta 60
cgttctcgag cgcgcgtaat acgactcact atagggcgaa ttgggtacgt tccatcatgg 120
agaaaaaaat cactggatat accaccgttg atatatccca atggcatcgt aaagaacatt 180
ttgaggcatt tcagtcagtt gctcaatgta cctataacca gaccgttcag ctggatatta 240
cggccttttt aaagaccgta aagaaaaata agcacaagtt ttatccggcc tttattcaca 300
ttcttgcccg cctgatgaat gctcatccgg aattccgtat ggcaatgaaa gacggtgagc 360
tggatgatag ggatagtggt cacccttggt acaccgtttt ccatgagcaa actgaaacgt 420
tttcatcgct ctggagtga taccacgacg atttccggca gtttctacac atatattcgc 480
aagatgtggc gtgttacggg gaaaacctgg cctatttccc taaaggggtt attgagaata 540
tgtttttcgt ctccagccaat cctgggtgga gtttcaccag ttttgattta aacgtggcca 600
atatggacaa cttcttcgcc ccggttttca ccatgggcaa atattatacg caaggcgaca 660
aggtgctgat gccgctggcg attcaggttc atcatgccgt ctgtgatggc ttccatgtcg 720
gcagaatgct taatgaatta caacagtact gcgatgagtg gcagggcggg gcgcgttaac 780
gagatcagct gaaaaatgat cttcttgaaa atttgcaggc cgtacgtgta ccggggcccc 840
cctcgactcg cgaaggagtc caccatgagt aaaggagaag aacttttcac tggagtgttc 900
ccaattcttg ttgaattaga tggatggtt aatgggcaca aattttctgt cagtggagag 960
ggatgaagtg atgcaacata cggaaaactt acccttaaat ttatttgac tactggaaaa 1020
ctacctgttc catggccaac acttgcactt actttcactt atgggtgttc atgcttttca 1080
agatacccag atcatatgaa acagcatgac tttttcaaga gtgccatgcc cgaagggtat 1140
gtacaggaaa gaactatatt tttcaaagat gacgggaact acaagacag tgctgaagtc 1200
aagtttgaag gtgataccct tgtaataaga atcgagttta aaggatttga ttttaaagaa 1260
gatggaaaca ttcttgga caaattggaa tacaactata actcacaaa tgtatacatc 1320
atggctgaca agcagaagaa cggaatcaag gccaaactta agaccgcgca caacatcgag 1380
gacggcggcg tgcagctggc cgaccactac cagcagaaca cccaatttgg cgatggccct 1440
gtccttttac cagacaacca ttacctgtcc acacaatctg ccctttcgaa agatcccaac 1500
gaaaagagag accacatggg cttcttgag tttgtaacag ctgctgggat tacacatggc 1560
atggatgaac tatacaaggg atcttcatga tctcagcaaa ctcttccttc ttaatccttc 1620
cagactcgaa gtcaattcgt gcatcaatcc gggccctaga caccatggcc tccaccatac 1680
tggaaattcc aactggtctt ctgtatgagc tgctagggaa gaatttctcg aatagggttc 1740
aacacttctg gtacatttgt tcactctcaa ggattccctt ttgactcgta ttgagaatgg 1800
aacggtttct cttagggatc caagagtgtg tagttgccac agcatcatat tccatgcttt 1860
tggctggacc atgggctggc attaccgcag cattgtttac agattcaatt tccttatgac 1920
tgacaaacgg gttcatggga ttacaaagtc ttccctgata gtcttcatcc attagtttcc 1980
atttcaggca aacttccggg atgtggagat tccgaatgtt gtacagggtt ggtecgccat 2040
ctgaaacca cagtctgcc tttgagcggg tctgtctccc cagcttcttt agctcgaatg 2100
acctcctcgt ttggatttgt gtgtctcccc tgtgacaccg gtatgtatat ctgtagtctt 2160
tgatgaataa ttggagagcc atttgggctg ttgccggtcc aagatcattg tttatcatgt 2220
tattctttat cactgttact ccaatgctca tatcagccga ttcattaatt cctgatactc 2280
caaagctggg caactccata ctaaaattgg ctacaaatcc atagcggtag aaaaagcttg 2340

EP 1 174 514 A1

	tgaattcgaa	tgttctgtc	ctatttatat	aggacttttt	cttgttcata	ttgatcccaa	2400
	ctagcttgca	ggttctgtag	aatctatcca	ctcccgcttg	tattccctca	tgatttggtg	2460
	cattcacgat	gagagcaaaa	tcacagagg	actgaagtc	atcccaccag	tatgtggttt	2520
5	tggtgtatct	cttttgccca	agattcagga	ttgagactcc	caacactgta	ctcagcatgt	2580
	tgaacatacc	catcatcatt	cccgggctta	atgagagctgt	gccgtctatt	atgagaggat	2640
	cgataggcct	agctaggtaa	agaaaaatac	ccttgtttct	actaataaac	cggcgcccca	2700
	aaatgccgac	toggagcgaa	agatatacct	cccccggggc	cgggaggtcg	cgtcacccgac	2760
	cacgcgcgcg	gcccaggcga	cgcgcgacac	ggacacctgt	ccccaaaaac	gccaccaTcg	2820
10	cagccacaca	cggagcgccc	ggggccctct	ggtcaacccc	aggacacacg	cgggagcagc	2880
	gccggggcgcg	ggagcgccctc	ccggccgccc	gtgccacacg	cagggggcgcg	gcccggtgtct	2940
	ccagagcggg	agccggaagc	attttcggcc	ggccctctct	acgaccggga	cacacgaggg	3000
	accgaaggcc	ggccaggcgc	gacctctcgg	gcccgcacgcg	cgtcaggga	gcgtctctccg	3060
	actccgcacg	gggactcgcc	agaaaggatc	gtgacctgca	ttaatgaatc	aggggataac	3120
15	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	aaaggccgcg	3180
	ttgctggcgt	ttttccatag	gctccgcccc	cctgacgagc	atcacaaaaa	tcgacgetca	3240
	agtccagaggt	ggcgaacccc	gacaggacta	taaagatacc	aggcggtttcc	ccctggaagc	3300
	tccctcgtgc	gctctcctgt	tccgacctg	ccgcttaccg	gatacctgtc	cgcctttctc	3360
	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	ggtatctcag	ttcgggtgta	3420
20	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccg	ccgctgcgcc	3480
	ttatccggta	actatcgtct	tgagtccaac	cgggtaagac	acgacttatc	gccactggca	3540
	gcagccactg	gtaacaggat	tagcagagcg	aggatatgtag	gcggtgctac	agagttcttg	3600
	aagtgggtggc	ctaactacgg	ctacactaga	aggacagtat	ttgggtatctg	cgtctgtctg	3660
	aagccagtta	ccttcggaaa	aagagttggt	agctcttgat	ccggcaaaaca	aaccaccgct	3720
25	ggtagcggtg	gtttttttgt	ttgcaagcag	cagattacgc	gcagaaaaaa	aggatctcaa	3780
	gaagatcctt	tgatcttttc	tacggggtct	gacgctcagt	ggaacgaaaa	ctcacgttaa	3840
	gggatttttg	tcagagagat	atcaaaaagg	atcttcacct	agatcctttt	aaattaaaaa	3900
	tgaagtttta	aatcaatcta	aagtatatat	gagtaaaact	ggtctgacag	ttaccaatgc	3960
	ttaatcagtg	aggcacctat	ctcagcgatc	tgtctatttc	gttcacccat	agttgcctga	4020
30	ctccccgtcg	gttagataac	tacgatacgg	gagggtttac	catctggccc	cagtgtcgca	4080
	atgataccgc	tgagaccacg	ctcaccggct	ccagatttat	cagcaataaa	ccagccagcc	4140
	ggaagggcgcg	agcgcagaag	tggctctgca	actttatccg	cctccatcca	gtctattaat	4200
	tggtgcccggg	aagctagagt	aagtagttcg	ccagttaata	gtttgcgcaa	cgttggtgcc	4260
	attgctacag	gcacgtgggt	gtcacgctcg	tcgtttggta	tggcttcatt	cagctccggt	4320
35	tcccaacgat	caaggcgagt	tacatgatcc	cccatggtgt	gcaaaaaagc	ggttagctcc	4380
	ttcggctctc	cgatcgttgt	cagaagtaag	ttggccgcag	tgttatcact	catgggtatg	4440
	gcagcaactgc	ataattctct	tactgtcatg	ccatccgtaa	gatgcttttc	tgtgactggt	4500
	gagtaactcaa	ccaagtcatt	ctgagaatag	tgtatgcggc	gaccgagttg	ctcttgcccc	4560
	gcgtcaacac	gggataatac	cgcgccacat	agcagaactt	taaaagtgtc	catcattgga	4620
40	aaacgttctt	cggggcgaaa	actctcaagg	atcttaccgc	tgttgagatc	cagttcgatg	4680
	taaccacactc	gtgcacccaa	ctgatcttca	gcacttttta	ctttcaccag	cgtttctggg	4740
	tgagcaaaaa	caggaaggca	aaatgccgca	aaaaagggaa	taagggcgac	acggaaatgt	4800
	tgaataactca	tactcttctc	ttttcaatat	tattgaagca	tttatcaggg	ttattgtctc	4860
45	atgagcggat	acatatttga	atgtatttag	aaaaataaac	aaaagagttt	gtagaaacgc	4920
	aaaaaggcca	tccgtcagga	tggccttctg	cttaatttga	tgcctggcag	tttatggcgg	4980
	gcgtcctgcc	cgccaccctc	cgggcggttg	cttcgcaacg	ttcaaatccg	ctcccgccgg	5040
	atttgtctca	ctcaggagag	cgttcaccga	caaacaacag	ataaaaacgaa	aggcccgatc	5100
	ttccgactga	gcctttcggt	ttatttgatg	ctcggcagtt	ccctactctc	gcatggggag	5160
50	acccacact	accactggcg	ctacggcggt	tcacttctga	gttcggcatg	gggtcaggtg	5220
	ggaccacccg	gctactgccg	ccaggcaaat	tctgttttat	cagaccgctt	ctgcgtttctg	5280
	atttaactctg	tatcaggctg	aaaatcttct	ctcatccgcc	aaaacagaag	ctagcggccg	5340
	atccccaaaa	aaaaaaaaaa	aaaaaaaaaa	aagagtcacg	agtggccccc	ccgttcgcgcg	5400
	ccggggggggg	gggggggggg	ggacactttc	ggacatctgg	tcgacctcca	gcacgggggg	5460
55	aaaaaaaaaa	aacaaagttt	cggccggagt	actggctcgac	ctccgaagtt	gggggggg	5517

EP 1 174 514 A1

<210> 24
 <211> 5699
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: pHL3236

<400> 24
 cctctcataa tagacggcac agcctcatta agccccgggaa tgatgatggg tatgttcaac 60
 atgctgagta cagtgttggg agtctcaatc ctgaatcttg ggcaaaagag atacaccaa 120
 accacatact ggtgggatgg acttcagtc tctgatgatt ttgctctcat cgtgaatgca 180
 ccaaatcatg aggggaatata agcgggagtg gatagattct acagaacctg caagctagtt 240
 gggatcaata tgagcaagaa aaagtcctat ataaatagga caggaacatt cgaattcaca 300
 agctttttct accgctatgg atttgtagcc aatttttagta tggagtgtgc cagctttgga 360
 gtatcaggaa ttaatgaatc ggctgatatg agcattggag taacagtgat aaagaataac 420
 atgataaaca atgactcttg accggcaaca gcccaaatgg ctctccaatt attcatcaag 480
 gactacagat atacataccg gtgtcacagg ggagacacac aaatccaaac gaggagggtca 540
 ttcgagctaa agaagctgtg ggagcagacc cgctcaaagg caggactgtt ggtttcagat 600
 ggcggaccac accgtgtaca cattcggaat ctccacatcc cggaagtgtt cctgaaatgg 660
 gaactaatgg atgaagacta tcagggaaga ctttgaatc ccatgaaccc gtttgtcagt 720
 cataaggaaa ttgaatctgt aaacaatgct gcggtaatgc cagcccatgg tccagccaaa 780
 agcatggaat atgatgctgt ggcaactaca cactcttggg tccctaagag aaaccgttcc 840
 attctcaata cgagtcaaag gggaatcctt gaggatgaac aaatgtacca gaagtgttgc 900
 aacctattcg agaaattctt ccctagcagc tcatacagaa gaccagtgtg aatttccagt 960
 atggtggagg ccattggtgtc tagggcccgg attgatgcac gaattgactt cgagtctgga 1020
 aggattaaga aggaagagtt tgctgagatc atgaagatcc cccgggctgc aggaattcga 1080
 tatcaagctt cgacgaattt ctgccattca tccgcttatt atcacttatt caggcgtagc 1140
 accaggcgtt taagggcacc aataactgcc ttaaaaaaat tacgccccgc cctgccactc 1200
 atcgagtcac tgttgaattt cattaagcat ctgcccagca tggaagccat cacaacggc 1260
 atgatgaacc tgatcgcca gcggcatcag caccctgtcg ccttgctgat aatatttgc 1320
 catggtgaaa acgggggcga agaagttgtc catattggcc acgtttaaat caaaactggt 1380
 gaaactcacc cagggtattg ctgagacgaa aaacatattc tcaataaacc ctttagggaa 1440
 ataggccagg ttttcaccgt aacacgccac atcttgcgaa tatatgtgta gaaactgccg 1500
 gaaatcgctg tggatttcac tccagagcga tgaaaacgtt tcagtgttgc catggaaaac 1560
 ggtgtaacaa ggggtgaacac tatcccatat caccagctca cctctttca ttgccatacg 1620
 gaattccgga tgagcattca tcaggcgggc aagaatgtga ataaaggccg gataaaactt 1680
 gtgcttattt ttctttacgg tctttaaaaa ggccgtaata tccagctgaa cggctctggt 1740
 ataggtacat tgagcaactg actgaaatgc ctcaaatgt tctttacgat gccattggga 1800
 tataatcaac gtggtatata cagtgtttt tttctccatt ttagcttctt tagctcctga 1860
 aaatctcgtc gaagcttatc gataccgtcg acctcgagg ggggcccgtt acggcctgca 1920
 aattttcaag aagatcattt ttacgtgat ctctgtatct agaccatgga gcttagtgat 1980
 ggtgatggtg atgggatccc ttgtatagtt catccatgcc atgtgtaatc ccagcagctg 2040
 ttacaaactc aagaaggacc atgtggtctc tcttttcgtt gggatctttc gaaagggcag 2100
 attgtgtgga caggtaatgg ttgtctggtg aaaggacagg gccatcgcca attgggggtg 2160
 tctgctggtg gtggtcggcc agctgcacgc cgccgtcctc gatgttgttg cgggtcttga 2220
 agttggcctt gattccgttc ttctgcttgt cagccatgat gtatacattg tgtgagttat 2280
 agttgtattc caatttgtgt ccaagaatgt ttccatcttc tttaaaatca atacctttta 2340
 actcgattct attaacaagg gtatcacctt caaacttgac ttcagcacgt gtctttagt 2400
 tcccgctatc tttgaaaaat atagttcttt cctgtacata accttcgggc atggcactct 2460
 tgaaaaagtc atgctgtttc atatgatctg ggtatcttga aaagcattga acaccataag 2520

EP 1 174 514 A1

	tgaaagtagt	gacaagtgtt	ggccatggaa	caggtagttt	tccagtagtg	caaataaatt	2580
	taagggttaag	ttttccgtat	gttgcacac	cttcaccctc	tccactgaca	gaaaatttgt	2640
	gcccatataac	atcaccatct	aattcaacaa	gaattgggac	aactccagtg	aaaagttctt	2700
5	ctcctttact	catggtggac	tccttcgcga	gtcagagggg	ggcccggtag	acgtacgcgc	2760
	tcgagaacgt	accttcgcgc	tcgcgatgtg	gatgtcactc	agtgagtgat	tatctaccct	2820
	gtttctactc	ccccccaact	tcggaggtcg	accagtactc	cgggcgaaac	tttgtttttt	2880
	ttttttcccc	cgatgctgga	ggtcgaccag	atgtccgaaa	gtgtccccc	ccccccccc	2940
	ccccggcgcg	gaacggcggg	gccactctgg	actctttttt	tttttttttt	tttttttttt	3000
10	ggggatcggc	cgtagcttc	tgttttggcg	gatgagagaa	gattttcagc	ctgatacaga	3060
	ttaaatcaga	acgcagaagc	ggtctgataa	aacagaattt	gcctggcggc	agtagcgcg	3120
	tgggtccacc	tgaccccatg	ccgaactcag	aagtgaacg	ccgtagcgcc	gatggtagtg	3180
	tggggtctcc	ccatgcgaga	gtaggggaact	gccaggcatc	aaataaaacg	aaaggctcag	3240
	tcgaaagact	gggcctttcg	ttttatctgt	tgtttgctcg	tgaacgctct	cctgagtagg	3300
15	acaaatccgc	cgggagcgga	tttgaacgtt	gcgaagcaac	ggcccggagg	gtggcgggca	3360
	ggacgcccgc	cataaactgc	caggcatcaa	attaagcaga	aggccatcct	gacggatggc	3420
	ctttttgctg	ttctacaaac	tcttttggtt	atttttctaa	atacattcaa	atatgtatcc	3480
	gctcatgaga	caataaccct	gataaatgct	tcaataatat	tgaaaaagga	agagtatgag	3540
	tattcaacat	ttccgtgtcg	cccttattcc	cttttttgcg	gcattttgcc	ttcctgtttt	3600
20	tgctcaccca	gaaacgctgg	tgaaagttaa	agatgctgaa	gatcagttgg	gtgcacgagt	3660
	gggttacatc	gaactggatc	tcaacagcgg	taagatcctt	gagagttttc	gccccgaaga	3720
	acgtttttcca	atgatgagca	cttttaaaagt	tctgctatgt	ggcgcggtat	tatcccggtg	3780
	tgacgcgggg	caagagcaac	tcggtcgccc	catacactat	tctcagaatg	acttggttga	3840
	gtactcacca	gtcacagaaa	agcatcttac	ggatggcatg	acagtaagag	aattatgcag	3900
25	tgctgccata	accatgagtg	ataacactgc	ggccaaactta	cttctgacaa	cgatcggagg	3960
	accgaaggag	ctaaccgctt	ttttgcacaa	catgggggat	catgtaactc	gccttgatcg	4020
	ttgggaaccg	gagctgaatg	aagccatacc	aaacgacgag	cgtgacacca	cgatgcctgt	4080
	agcaatggca	acaacgttgc	gcaaacattt	aactggcgaa	ctacttactc	tagcttcccc	4140
	gcaacaatta	atagactgga	tgagggcgga	taaagtgcga	ggaccacttc	tgcgctccgc	4200
30	ccctccggct	ggctgggtta	ttgctgataa	atctggagcc	ggtgagcgtg	ggtctcgcgg	4260
	tatcattgca	gcactggggc	cagatggtaa	gccctcccg	atcgtagtta	tctacacgac	4320
	ggggagtcag	gcaactatgg	atgaacgaaa	tagacagatc	gctgagatag	gtgcctcact	4380
	gattaagcat	tggttaactgt	cagaccaagt	ttactcata	atactttaga	ttgatttaaa	4440
	acttcatttt	taattttaaaa	ggatctaggt	gaagatcctt	tttgataatc	tcatgaccaa	4500
35	aatcccttaa	cgtgagtttt	cggtccactg	agcgtcagac	cccgtagaaa	agatcaaagg	4560
	atcttcttga	gatccttttt	ttctgcgcgt	aatctgctgc	ttgcaaacia	aaaaaccacc	4620
	gctaccagcg	gtggtttggt	tgcgggatca	agagctacca	actctttttc	cgaaggtaac	4680
	tggttccagc	agagcgcaga	taccaaatac	tgctcttcta	gtgtagccgt	agttaggcca	4740
	ccacttcaag	aactctgtag	caccgcctac	atacctcgct	ctgctaatac	tgttaccagt	4800
40	ggctgctgcc	agtggcgata	agtcgtgtct	taccgggttg	gactcaagac	gatagttacc	4860
	ggataaggcg	cagcggtcgg	gctgaacggg	gggttcgtgc	acacagccca	gcttggagcg	4920
	aacgacctac	accgaactga	gatacctaca	gcgtgagcta	tgagaaagcg	ccacgcttcc	4980
	cgaaggagga	aaggcggaca	ggtatccggt	aagcggcagg	gtcggaaacg	gagagcgcac	5040
	gaggagagctt	ccagggggaa	acgcctggta	tctttatagt	cctgtcgggt	ttcgccacct	5100
45	ctgacttgag	cgctgatttt	tgatgatgtc	gtcagggggg	cggagcctat	ggaaaaacgc	5160
	cagcaacgcg	gccttttttac	ggttctctgg	cttttgctgg	ccttttgctc	acatgttctt	5220
	tcttgctgta	tcccttgatt	cattaatgca	ggtcacgata	ctttctggcg	agtcctcggt	5280
	cggagtcgga	tgagcgcgcc	tgagggccga	gaggtcgcgc	ctggccggcc	ctggccggcc	5340
	ttcggccctc	cgtgtgtccc	ggtcgtagga	ggggccggcc	gaaaatgctt	ccggctcccg	5400
50	ctctggagac	acgggcgggc	ccctgcgctg	tggcacgggc	ggccgggagg	gcgtccccgg	5460
	cccggcgctg	ctcccgctg	tgctctgggg	ttgaccagag	ggccccgggc	gctccgtgtg	5520
	tggctgcgat	ggtggcggtt	ttggggacag	gtgtccgtgt	cgcgcgtcgc	ctgggcgggc	5580
	ggcgtggctg	gtgacgcgac	ctcccgcccc	cgggggaggt	atatctttcg	ctccgagtcg	5640
55	gcattttggg	ccgccggggt	attagtagaa	acaagggtat	ttttctttac	ctagctagg	5699

Claims

1. A recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus and which comprises at least one viral RNA segment being a bicistronic RNA molecule coding for two genes in tandem arrangement (tandem RNA segment), in said tandem RNA segment one of the standard viral genes being in covalent junction with a foreign, recombinant gene and said tandem RNA segment having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region.
2. The recombinant influenza virus of claim 1, wherein the tandem RNA segment contains one of the standard viral genes in distal mRNA position behind a foreign, recombinant gene in proximal position, or vice versa, both in antisense orientation with regard to the viral RNA as present within the virus.
3. The recombinant influenza virus of claim 1 or 2, wherein at least one of the regular viral RNA segments is replaced by a tandem RNA segment, preferably the replaced regular viral RNA segment is selected from the neuraminidase segment, hemagglutinin segment and NS segment.
4. The recombinant influenza virus of claims 1 to 3, wherein the splice donor and splice acceptor signals are selected from sequences as present in influenza WSN segment 7 and 8 or other partially effective splice reactin substrates.
5. The recombinant influenza virus of claim 4, wherein the splice donor and splice acceptor signals are selected from sequences as present in influenza WSN segment 7.
6. The recombinant influenza virus according to claims 1 to 5, wherein one or more of the regular viral RNA segments, differing from said at least one tandem RNA segment, comprises a vRNA encoding a foreign gene which may or may not be in covalent connection to one of the viral genes, and preferably one or more of the regular viral RNA segments has (have) been deleted and replaced by a tandem vRNA encoding in addition a foreign gene.
7. The recombinant influenza virus according to claims 1 to 6, in which the terminal viral RNA sequences of one or more of the regular segments and/or of the at least one tandem RNA segment, which are active as the promoter signal, have been modified by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence.
8. The recombinant influenza virus of claim 7, wherein the 12 nucleotide conserved influenza 3' terminal sequence has been modified by replacement of one to three nucleotides occurring in said sequence at positions 3, 5 and 8 relative to the 3' end by other nucleotides, and/or wherein the 13 nucleotide conserved influenza 5' terminal sequence has been modified by replacement of one or two nucleotides occurring in said sequence at positions 3 and 8 by other nucleotides.
9. The recombinant influenza virus of claim 8, wherein the replacements in the 3' terminal nucleotide sequence comprises the modifications G3A and C8U.
10. The recombinant influenza virus of claim 9, wherein the replacements in the 3' terminal nucleotide sequence comprises the modifications G3A, U5C and C8U, or G3C, U5C and C8G.
11. The recombinant influenza virus of claim 10, which comprises a 3' terminal nucleotide sequence of (5')-CCUGU-UUCUACU-3'.
12. The recombinant influenza virus according to claims 7 to 12, wherein the 5' terminal nucleotide sequence comprises the modifications U3A and A8U resulting in a 5'-terminal sequence of 5'-AGAAGAAUCAAGG.
13. The recombinant influenza virus according to claims 1 to 12, which is a recombinant influenza A virus.
14. The recombinant influenza virus according to claims 1 to 13, in which the foreign gene(s) in the tandem RNA segment code for proteins and/or glycoproteins which are secreted from cells infected with the recombinant virus.
15. The recombinant influenza virus according to claims 1 to 13, in which the foreign gene(s) in the tandem RNA segment code for proteins or artificial polypeptides designed to support an efficient presentation of inherent

epitopes at the surface of infected cells, for stimulation of a B cell and/or T cell response.

16. The recombinant influenza virus according to claims 1 to 13, in which the foreign gene(s) in the tandem RNA segment is a nucleotide sequence causing viral attenuation.

17. The recombinant influenza virus of claim 16, wherein the foreign gene is coding for part of or for the entire viral neuraminidase gene in antisense orientation.

18. The recombinant influenza virus of claim 17, wherein the neuraminidase gene in antisense orientation is attached to the hemagglutinin vRNA segment, and optionally another gene or reporter gene is encoded in a second tandem vRNA, preferably in conjunction with NS2.

19. A method for the production of recombinant influenza viruses as defined in claims 1 to 18 comprising

- (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense or in sense tandem design,
- (b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in the corresponding viral RNA segment, and
- (c) thereafter selective vRNA inactivation through ribozyme cleavage.

20. A pharmaceutical composition comprising a recombinant influenza virus according to claims 1 to 18, preferably a recombinant influenza virus of claims 16 to 18.

21. Use of a recombinant influenza virus according to claims 1 to 18, preferably a recombinant influenza virus of claims 16 to 18, for preparing a medicament for vaccination purposes.

22. The use according to claim 21, wherein the medicament

- (a) is suitable against influenza and/or against other infections;
- (b) is present in form of inactivated preparations; and/or
- (c) is present in form of live recombinant viruses.

23. Use of a recombinant influenza virus according to claims 1 to 18 for preparing agents for somatic gene therapy.

24. Use of a recombinant influenza virus according to claims 1 to 18 for preparing agents, for transfer and expression of foreign genes into cells infected by such viruses.

25. Use of a recombinant influenza virus according to claims 1 to 18 for preparing agents for transfer and expression of RNA molecules into cells infected by such viruses.

26. The use of claim 24, wherein the RNA molecules to be expressed are antisense sequences or double-strand sequences relative to the target cell cellular mRNA molecules, and/or the agent is suitable for sequence-specific gene silencing, preferably by antisense RNA or RNA interference mechanisms.

27. The use according to claims 23 to 26, wherein the agents are applicable in *ex vivo* and *in vivo* application schemes.

28. A method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus according to claims 1 to 19 as expression vector.

29. The method of claim 28, wherein the production is performed in cell culture cells or in fertilized chicken eggs.

30. A method for preventing and/or treating influenza which comprises administering an effective amount of a recombinant influenza virus according to claims 1 to 18, preferably of a recombinant influenza virus according to claims 16 to 18, to the mammal to be treated.

31. A method for somatic gene therapy, which method comprises subjecting the organism to be treated with a recombinant influenza virus according to claims 1 to 18.

32. A method for transfer and expression of foreign genes into cells, and for transfer and expression of RNA molecules

EP 1 174 514 A1

into cells, which method comprises infecting the cells with a recombinant influenza virus according to claims 1 to 18.

33. Use of a recombinant influenza virus according to claims 1 to 18 for preparing agents for immunotherapy, preferably for autologous immunotherapy.

5

34. A method for an immunotherapy which comprises *ex vivo* infection of immune cells, preferably dendritic cells, with a recombinant influenza virus according to claims 1 to 18, and introduction of the transduced cells into the patient.

35. A method for the induction of antibodies which comprises utilizing a recombinant influenza virus according to claims 1 to 18 as an immunogen.

10

15

20

25

30

35

40

45

50

55

FIG. 1




FIG. 2




FIG. 3





FIG. 4

A



B




FIG. 5






FIG. 7





FIG. 8





European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 00 11 5626 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL7)
A	FLICK & HOBOM: "Transient bicistronic vRNA segments for indirect selection of recombinant influenza viruses." VIROLOGY. 1999 SEP 15;262(1):93-103., XP000986453 * the whole document *	1,7,11, 13,15,18	C12N15/86 A61K48/00 C12N7/04 A61K39/145
A	DE 197 09 512 A (HOBOM GERD PROF DR DR) 10 September 1998 (1998-09-10) * the whole document *	1,4, 13-15, 20-35	
A	HOFFMANN E ET AL.: "'Ambisense" approach for the generation of influenza A virus: vRNA and mRNA synthesis from one template." VIROLOGY. 2000 FEB 15;267(2):310-7., XP000986499 * the whole document *	1	
			TECHNICAL FIELDS SEARCHED (InCL7)
			C12N
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		15 February 2001	Chambonnet, F
CATEGORY OF CITED DOCUMENTS			
<p>X: particularly relevant if taken alone</p> <p>Y: particularly relevant if combined with another document of the same category</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p>		<p>T: theory or principle underlying the invention</p> <p>E: earlier patent document, but published on, or after the filing date</p> <p>D: document cited in the application</p> <p>L: document cited for other reasons</p> <p>&: member of the same patent family, corresponding document</p>	

EPO FORM 1503 (03.02.92) (954/07)



European Patent
Office

**INCOMPLETE SEARCH
SHEET C**

Application Number
EP 00 11 5626

As far as claims 28, 29, 32 and 35 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), as well as claims 30, 31 and 34 that are directed to such a method, the search has been carried out and based on the alleged effects of the compound/composition.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 00 11 5626

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	NEUMANN G ET AL: "Plasmid-driven formation of influenza virus-like particles" JOURNAL OF VIROLOGY, THE AMERICAN SOCIETY FOR MICROBIOLOGY, US, vol. 74, no. 1, January 2000 (2000-01), pages 547-551, XP002140119 ISSN: 0022-538X * the whole document *	1	
A	HOFFMANN, E ET AL.: "A DNA transfection system for generation of influenza A virus from eight plasmids." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol. 97, no. 11, 23 May 2000 (2000-05-23), pages 6108-6013, XP000982038 NATIONAL ACADEMY OF SCIENCE. WASHINGTON., US ISSN: 0027-8424 * the whole document *	1	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
T,D	WO 00 53786 A (AZZEH MAYSA ;HOBOM GERD (DE); MENKE ANETTE (DE); ARTEMIS PHARMACEU) 14 September 2000 (2000-09-14) * the whole document *	1	

EP 00 11 5626 (P44C18)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 11 5626

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

15-02-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19709512 A	10-09-1998	NONE	
WO 0053786 A	14-09-2000	EP 1035209 A AU 3427100 A	13-09-2000 28-09-2000

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82